


Question	Scheme	Marks	AOs
2(a)		B1	1.1b
		(1)	
(b)	<p>Using <i>total</i> area = 15000 to set up an <i>equation</i> in <b>one unknown</b>  Or they may use <i>suvat</i> on one or more sections (but must still be considering <i>all</i> sections)</p> <p>Allow an attempt at a clear explicit verification using <math>t = 40</math>  e.g. the following would score M1A1A1*:  <math>4 \times 40 = 160</math> then <math>700 - 40 - 160 = 500</math>  <math>\frac{(700 + 500)}{2} \times 25 = 15000 = 15 \text{ km}</math>  Withhold A1* if they don't include <math>= 15 \text{ km}</math></p> <p><b>N.B.</b> M0 if a single <i>suvat</i> formula is used for the whole journey.</p>	M1	3.4
	$\frac{1}{2}(700 + 700 - t - 4t) \times 25 = 15000$ <p><b>OR</b> <math>\frac{1}{2} \times 25 \times t + 25(700 - t - 4t) + \frac{1}{2} \times 25 \times 4t = 15000</math></p>	A1	1.1b
	$t = 40 \text{ (s)*}$	A1*	1.1b
		(3)	
(c)	0.63 or 0.625 or $\frac{5}{8}$ oe (m s <sup>-2</sup> ) isw	B1	1.1b/ (2.2a)
		(1)	
(d)	<p>Complete method to find the speed or velocity at <math>t = 572</math>  e.g. <math>\pm \left( 25 - \left( 32 \times \frac{5}{32} \right) \right)</math> or <math>\pm \left( 128 \times \frac{5}{32} \right)</math> oe</p>	M1	3.1b
	20 (m s <sup>-1</sup> )	A1	1.1b
		(2)	
(e)	<p>e.g. (the train) cannot instantaneously change acceleration, (the train) won't move with <u>constant</u> acceleration , (the train) won't move with <u>constant</u> speed  Allow negatives of these:</p>	B1	3.5b

	<p>e.g. (The train) moving at constant speed, or just ‘constant speed’ or ‘constant acceleration’ (is a limitation of the model)</p> <p>Must be a limitation of the model, so friction or air resistance or size of train is B0.</p> <p><b>N.B.</b> Ignore incorrect <b>reasons</b> following a correct answer.</p>		
		(1)	

(8 marks)

Notes:

2a	B1	<p>Overall shape of graph, starting at the origin, with deceleration phase <i>longer</i> than the acceleration phase if nothing on the <math>t</math>-axis but ignore the relative lengths and allow if <math>t</math> (or 40) and <math>4t</math> (or 160) are clearly and correctly marked.</p> <p>Ignore incorrect figs on the axes.</p> <p>This mark can be earned if the graph appears anywhere in qu 2.</p>
2b	M1	<p>Need <i>all</i> sections to be included, with <u>correct structure for each section</u>, with <math>\frac{1}{2}</math> 's where appropriate.</p> <p><u>Allow = 15 or 150 or 1500 etc instead of 15000</u></p>
	A1	A correct equation <b>in their <math>t</math> only, seen or implied</b> (or with $t = 40$ for verification)
	A1*	cso. At least one line of working with brackets removed and $t$ 's collected, or equivalent
2c	B1	cao
2d	M1	Any complete method, must have correct figs, but condone sign errors
	A1	cao. Must be positive and exact i.e must not come from rounding.
2e	B1	Any appropriate limitation <u>of the model</u> . B0 if any incorrect extra <b>answers</b> .