

Question	Scheme	Marks	AOs
4(a)	The horizontal component of $T$ acts to the left and since the <b>only</b> other horizontal force is friction, it must act to the right oe	B1	2.4
		(1)	
4(b)	Take moments about A or any other complete method to obtain <b>an equation in <math>T</math>, <math>M</math> and <math>\theta</math> only</b> . (see possible equations below that they may use)	M1	3.1b
	$T.2a = Mga \cos \theta + 2Mg \times 1.5a \cos \theta$ (A0 if $a$ 's missing)	A1	1.1b
	Other possible equations but $F$ and $R$ would need to be eliminated. $(\nwarrow), R \cos \theta + T = F \sin \theta + Mg \cos \theta + 2Mg \cos \theta$ $(\nearrow), R \sin \theta + F \cos \theta = Mg \sin \theta + 2Mg \sin \theta$ $(\rightarrow), F = T \sin \theta$ M(B), $R.2a \cos \theta = Mga \cos \theta + 2Mg \times 0.5a \cos \theta + F.2a \sin \theta$ M(G), $Fa \sin \theta + Ta = Ra \cos \theta + 2Mg \times 0.5a \cos \theta$ M(C), $R \times 1.5a \cos \theta = T \times 0.5a + Mg \times 0.5a \cos \theta + F \times 1.5a \sin \theta$		
	$T = 2Mg \cos \theta^*$	A1*	1.1b
		(3)	
4(c)	e.g. Resolve vertically	M1	3.4
	$(\uparrow), R + T \cos \theta = Mg + 2Mg$	A1	1.1b
	$R = \frac{57Mg}{25}^*$	A1*	1.1b
		(3)	
	Other possible equations but $F$ would need to be eliminated. $(\nwarrow), R \cos \theta + T = F \sin \theta + Mg \cos \theta + 2Mg \cos \theta$ $(\nearrow), R \sin \theta + F \cos \theta = Mg \sin \theta + 2Mg \sin \theta$ $(\rightarrow), F = T \sin \theta$ M(B), $R.2a \cos \theta = Mga \cos \theta + 2Mg \times 0.5a \cos \theta + F.2a \sin \theta$ M(G), $Fa \sin \theta + Ta = Ra \cos \theta + 2Mg \times 0.5a \cos \theta$ M(C), $R \times 1.5a \cos \theta = T \times 0.5a + Mg \times 0.5a \cos \theta + F \times 1.5a \sin \theta$		
4(d)	Find an equation containing $F$ e.g. Resolve horizontally	M1	3.4
	$(\rightarrow), F = T \sin \theta$	A1	1.1b
	Other possible equations		

	$(\nwarrow), R \cos \theta + T = F \sin \theta + Mg \cos \theta + 2Mg \cos \theta$ $(\nearrow), R \sin \theta + F \cos \theta = Mg \sin \theta + 2Mg \sin \theta$ $(\rightarrow), F = T \sin \theta$ $M(B), R.2a \cos \theta = Mga \cos \theta + 2Mg \times 0.5a \cos \theta + F.2a \sin \theta$ $M(G), Fa \sin \theta + Ta = Ra \cos \theta + 2Mg \times 0.5a \cos \theta$ $M(C), R \times 1.5a \cos \theta = T \times 0.5a + Mg \times 0.5a \cos \theta + F \times 1.5a \sin \theta$		
	$F = \mu R$ used i.e. both $F$ and $R$ are substituted.	M1	3.1b
	$\mu = \frac{8}{19} *$	A1*	2.2a
		(4)	

(11 marks)

Notes:

4a	B1	Any equivalent explanation
4b	M1	Correct no. of terms, dimensionally correct, condone sin/cos confusion and sign errors
	A1	Correct equation, trig does not need to be substituted (Allow: $T.2a = Mga \cos \theta + 3Mga \cos \theta$ )
	A1*	Given answer correctly obtained with <u>no wrong working seen</u> . Allow $2Mg \cos \theta = T$ But not $T = 2 \cos \theta Mg$
4c	M1	For an equation in $R, M, T$ and $\theta$ <b>only</b> Correct no. of terms, dimensionally correct, condone sin/cos confusion and sign errors, each term that needs to be resolved must be resolved
	A1	Correct equation, $T$ and trig do not need to be substituted
	A1*	Given answer correctly obtained with <u>no wrong working seen</u>
4d	M1	For any equation with $F$ in it Correct no. of terms, dimensionally correct, condone sin/cos confusion and sign errors, each term that needs to be resolved must be resolved
	A1	Correct equation, trig does not need to be substituted
	M1	Must be used i.e M0 if merely quoting it.
	A1*	Given answer correctly obtained with <u>no wrong working seen</u>